Appendix A

l. - t r r t r 100,000 f r - t 20] 2020, r t r t f r ff r t 2020 20] (t 5% f t w), v r

People	7.4 (7.1, 7.8)	7.2 (6.8, 7.5)	0.97 (0.91, 1.03)
Male	9.7 (9.1, 10.2)	9.5 (9.0, 10.1)	0.98 (0.91, 1.07)
Female	5.2 (4.8, 5.6)	4.9 (4.5, 5.3)	0.94 (0.85, 1.05)

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2019 and 2020 are preliminary and thus are subject to further revision. 95% confidence intervals for the age-standardised rate and rate ratio are shown in brackets. Please refer to our methods

2. r r t r 100,000 f r - t 20] 2020, r t r t f r ff r t 2020 20] (t 5% f t w), r

15-24	3.5 (2.9, 4.2)	3.7 (3.1, 4.5)	1.06 (0.81, 1.38)
25-34	8.0 (7.1, 8.9)		

Major Cities	15-24	3.9 (3.1, 4.7)	4.1 (3.3, 5.0)	1.05 (0.79, 1.41)
Major Cities	25-34	7.7 (6.7, 8.7)	7.7 (6.7, 8.8)	1.00 (0.83, 1.21)
Major Cities	35-44	13 (12, 15)	13 (12, 15)	1.00 (0.86, 1.16)
Major Cities	45-54	15 (13, 17)	13 (12, 15)	0.88 (0.75, 1.03)
Major Cities	55-64	9.6 (8.2, 11.0)	10 (9, 11)	1.04 (0.85, 1.28)
Major Cities	65-74	5.3 (4.2, 6.6)	4.7 (3.6, 5.9)	0.88 (0.63, 1.23)
Major Cities	75-84	5.3 (3.8, 7.0)	5.1 (3.7, 6.8)	0.95 (0.61, 1.48)
Major Cities	85+	7.8 (4.9, 10.9)	10 (7, 14)	1.32 (0.79, 2.24)
Regional and Remote				

Antipsychotics & neuroleptics	1.5 (1.3, 1.6)	1.2 (1.1, 1.4)	0.84 (0.72, 0.98)
Cannabinoids	0.85 (0.74, 0.98)	0.44 (0.36, 0.54)	0.52 (0.41, 0.66)
Cocaine	0.35 (0.28, 0.43)	0.35 (0.28, 0.43)	0.99 (0.73, 1.34)
Nonopioid analgesics	1.2 (1.0, 1.3)	0.77 (0.66, 0.88)	0.65 (0.55, 0.78)
Opioids	4.6 (4.3, 4.8)	4.3 (4.1, 4.6)	0.95 (0.87, 1.03)

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2019 and 2020 are preliminary and thus are subject to further revision. 95% confidence intervals for the age-standardised rate and rate ratio are shown in brackets. Please refer to our methods document on 'Presentation of results' for interpretation of rate ratios. Please also refer to our methods document on 'Data source' and 'Coding of deaths' for details on the data used.

11. - t r r t r 100,000 f r - t 20] 2020, r t r t f r ff r t 2020 20] (t 5% f t w), t t(t t t t) r t v v

Unintentional	Alcohol	0.89 (0.78, 1.02)	0.83 (0.72, 0.95)	0.93 (0.77, 1.12)
Unintentional	Amphetamines	1.7 (1.5, 1.8)	1.8 (1.6, 2.0)	1.09 (0.95, 1.25)
Unintentional	Antidepressants	1.4 (1.2, 1.5)	1.3 (1.1, 1.4)	0.93 (0.80, 1.08)
Unintentional				

the age-standardised rate and rate ratio are shown in brackets. Please refer to our methods document on 'Presentation of results' for interpretation of rate ratios. Please also refer to our methods document on 'Data source' and 'Coding of deaths' for details on the data used.

12. - t r r t r 100,000 f r - t v v t r 20] 2020, r t r t f r ff r t 2020 20] (t 5% f t w),

Amphetamines	Female	1.0 (0.9, 1.2)	1.2 (1.1, 1.5)	1.19 (0.94, 1.51)
Amphetamines	Male	2.9 (2.6, 3.2)	3.1 (2.8, 3.4)	1.07 (0.92, 1.23)
Amphetamines	People	2.0 (1.8, 2.1)	2.1 (2.0, 2.3)	1.10 (0.97, 1.25)
Cocaine	Female	0.12 (0.07, 0.20)	NA (NA, NA)	NA (NA, NA)
Cocaine	Male	0.58 (0.45, 0.73)	0.63 (0.50, 0.79)	1.10 (0.80, 1.52)
Cocaine	People	0.35 (0.28, 0.43)	0.35 (0.28, 0.43)	0.99 (0.73, 1.34)

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2019 and 2020 are preliminary and thus are subject to further revision. 95% confidence intervals for the age-standardised rate and rate ratio are shown in brackets. Please refer to our methods document on 'Presentation of results' for interpretation of rate ratios. Please also refer to our methods document on 'Data source' and 'Coding of deaths' for details on the data used.

13. r r t r 100,000 f r - t v v t 20] 2020, r t r t f r ff r t 2020 20] (t 5% f t w),

	1.2 (0.8, 1.6)	1.2 (0.9, 1.7)	1.07 (0.67, 1.71)

results' for interpretation of rate ratios. Please also refer to our methods document on 'Data source' and 'Coding of deaths' for details on the data used.

15. r r t r 100,000 f - t 20 2020, r t r t f r ff r t 2020 20 (t 5% f t w),

15-24	2.1 (1.7, 2.7)	1.7 (1.3, 2.3)	0.82 (0.57, 1.19)
25-34	4.6 (3.9, 5.3)	5.0 (4.3, 5.8)	1.10 (0.89, 1.36)
35-44	9.5 (8.5, 10.6)	9.0 (8.0, 10.1)	0.94 (0.81, 1.11)
45-54	9.7 (8.7, 10.9)	8.4 (7.4, 9.4)	0.86 (0.73, 1.01)
55-64	5.9 (5.0, 6.8)	5.3 (4.5, 6.2)	0.91 (0.73, 1.14)
65-74	1.9 (1.4, 2.6)	2.3 (1.8, 3.0)	1.21 (0.80, 1.84)
75-84	1.5 (0.9, 2.3)	1.4 (0.8, 2.2)	0.95 (0.47, 1.94)
85+	1.6 (0.7, 3.1)	1.9 (0.9, 3.5)	1.22 (0.43, 3.55)

Note: Deaths where conditions related to alcohol or tobacco comprised the underlying cause of death are not captured here. Causes of death data for 2019 and 2020 are preliminary and thus are subject to further revision. 95% confidence intervals for the crude rate and rate ratio are shown in brackets. Please refer to our methods document on 'Presentation of results' for interpretation of rate ratios. The estimates for the 0-14 years age group are not presented due to sensitivity of the data. Please also refer to our methods document on 'Data source' and 'Coding of deaths' for details on the data used.

1 . -