ISSN 2982-0782

DOI https://doi.org/10.26190/unsworks/30193

Copyright ©NDARC, UNSW SYDNEY 2024

This report was prepared by researchers from the National Drug and Alcohol Research Centre for the Drug Trends program. The Drug Trends program is coordinated by the National Drug and Alcohol Research Centre, UNSW Sydney and undertaken in partnership with the Burnet Institute, National Drug Research Institute, University of Queensland, and University of Tasmania

This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use or use within your organisation. All other rights are reserved. Requests and

Last updated:17 June 2024

Trends in

List of Figures

Figure 1. Age-

Panel

Sociodemographic Characteristics of Drug-Related Hospitalisations

Sex

In 2021-22, drug-related hospitalisations were equally distributed among males and females (0 g 0 G [( )] TJ ET Q q 0.000009

Trends in Drug-

Drug-Related Hospitalisations by Drug

In 2021-22, the largest number of drug-related hospitalisations identified by principal diagnosis among the Australian population were attributable to amphetamine-type stimulants (mostly comprising methamphetamine), followed by antiepileptic, sedative-hypnotic and antiparkinsonism drugs (e.g., benzodiazepines; GHB), non-opio

Figure 8. Age-standardised rate per 100,000 people of drug -related hospitalisations among the Australian population, by drug identified in the principal diagnosis, 2002-03 to 2021-22.



Note: Please also refer to our methods document on 'Scope of the data' and 'Coding of hospitalisations' for specifications of data selected and all exclusions

Drug Type and Remoteness Area of Usual Residence

In 2021-22, amphetamine-type stimulants was the most common drug class identified as the principal diagnosis for drug -related hospitalisations in all areas, except for remote and very remote areaswhere cannabinoid-related hospitalisations remained the most common (Figure 9).

A detailed description of the distribution of selected drug -related hospitalisation rates across remoteness areas can be found in the subsequent sections.

Figure 9. Age-standardised rate per 100,000 people of drug -related hospitalisations, by drug type identified in the principal diagnosis and remoteness area, among the Australian population , 2018-19 to 2021-22.



Note: Age-standardised rates are not shown for remote and very remote areas for some drug types because the number of hospitalisations was less than or equal to 10. Please refer to our methods document for details.

Opioid-Related Hospitalisations

The following findings describe opioid-related hospitalisations due to illicit opioids (e.g., heroin), as well as opioids typically

Principal Diagnosis

In 2021-22, nearly half (49%) of all opioid-related hospitalisations were due to opioid poisoning (2,776 hospitalisations; 10 per 100,000 people), with slightly more of these poisonings determined to be intentional (45%; 1,242 hospitalisations; 4.8 per 100,000 people) than

Trends in Drug-Related Hospitalisations in

Amphetamine-Type Stimulant-Related Hospitalisations

The following findings describe hospitalisations with the principal diagnosis identified as amphetamine-type stimulant. This includes methamphetamines, 3,4-P H W K \ O H Q H G L R [\ P H W K D P S K H W D P L Q H 0 ' 0 $ ¶ H F V W D V (e.g., dexamphetamine), and other stimulants (e.g., caffeine).

In 2021-22, there were 32,010 hospitalisations where amphetamine-type stimulants were recorded as a principal or other diagnosis, among which 11,647 hospitalisations specifically identified amphetamine-type stimulants as a principal diagnosis. The latter figure translates to a population rate of 48 hospitalisations per 100,000 people.

Trend since 2002-03

- From 2002-03, the rate of hospitalisations related to amphetamine-type stimulants increased from 17 hospitalisations per 100,000 people to a peak of 70 hospitalisations per 100,000 people in 2019-20, before subsequently decreasing to 60 hospitalisations per 100,000 people in 2020-21 (Figure 15).
- In 2021-22 the rate recorded a further annual decrease of 20% (Table A1, Appendix).

Sex

In 2021-22, hospitalisations related to amphetamine-type stimulants were about 1.5 times as prevalent in males as in females, constituting 63% of all amphetamine-type stimulant-related hospitalisations, with a rate of 60 hospitalisations per 100,000 people for males compared to 36 hospitalisations per 100,000 people for females (Figure 15).

Trend since 2002-03

- The rate of h

- Of all age groups, the 50-59 age group recorded the biggest decrease from 2020-21 to 2021-22, which followed a 19-fold increase observed between 2002-03 and 2020-21 (Table A2, Appendix).
- The rates in the 60 and over age group stayed low and stable throughout monitoring.

Sex and Age

Trends in amphetamine-type related hospitalisations for males and females by age group follow a similar pattern as described above (Figure

Figure 2(e)] TJ ET Q q 0.00000912 0 612 792 re W* n BT /F4 9 Tf 0 0 0 rg 612 7

Principal Diagnosis

Over the 20-year monitoring period, amphetamine-type stimulant-related hospitalisations mostly comprised a principal diagnosis of mental and behavioural disorder due to substance use (92% in 2021-22). Among those who received a principal diagnosis of mental and behavioural disorder due to use of amphetamine-type stimulants in 2021-22 (10,699 hospitalisations; 44 per 100,000 people), drug-induced psychotic disorder was the most common reason for hospitalisation (49%; 5,237 hospitalisations; 21 per 100,000 people), followed by dependence (23%; 2,444 hospitalisations; 9.9 per 100,000 people) and acute intoxication (14%; 1,464 hospitalisations; 6.0 per 100,000 people), all of which declined in 2021-22 compared to 2020-21 (Figure 17) (Table A14, Appendix).

Please refer to the visualisation tool for trends over time by diagnosis type, although it is important to note changes over time may partly reflect changes in coding practices.

Figure 17. Age-standardised rate per 100,000 people of amphetamine-type stimulant-related hospitalisations among the Australian population, by principal diagnosis of mental and behavioural disorder due to substance use (A) and external cause of poisoning (B), 2002-03 to 2021-22.

Note: g 0 Gil Q qt5

7

Cannabinoid-Related Hospitalisations

In 2021-22, there were 40,244 hospitalisations with a cannabinoid-related diagnosis (including cannabis and

Figure 19. Age-

Figure 20. Crude rate per 100,000 people of cannabinoid-related hospitalisations among the female (A), male (B) and total (C) Australian population, by age group, 2002-03 to 2021-22.

Note: Given the small numbers, the age groups 60-69 years, and 70 years and over are combined into the 60 years and over age group. Numbers for the 50-59 years, and the 60 years and over age groups in the earlier years are small and thus rates are suppressed to protect confidentiality. The rates for the 0-9 years age group are not presented due to sensitivity of the data.

Remoteness Area of Usual Residence

- The highest rate of cannabinoid-related hospitalisations was observed in remote and very remote areas, with a rate similar to that observed in 2020-21 (70 versus 72 hospitalisations per 100,000 people in 2020-21).
- This rate was two times the rate in outer regional Australia (35 hospitalisations per 100,000 people), and nearly three times the rate in inner regional and in major city areas (26 and 25 hospitalisations per 100,000 people, respectively) (Figure 9).
- From 2020-21 to 2021-22, the hospitalisation rate declined in outer regional areas, dropping from 43 to 35 hospitalisations per 100,000 people. Other rates in other remoteness areas remained similar to those recorded in 2020-21.

Principal Diagnosis

Over the period of monitoring, cannabinoid-related hospitalisations mostly comprised a principal diag4 Tm 7S6-8(s)] TJ ET Q q

Principal Diagnosis

10

Hospitalisations Related to Other Drugs

There was a decline in the rate of hospitalisations with a principal diagnosis related to antiepileptic, sedative-hypnotic and antiparkinsonism drugs, from 51 hospitalisations per 100,000 people in 2002-03 to 31 hospitalisations per 100,000 people in 2021-22 (Figure 8). In 2021-22, nearly half of the hospitalisations related to antiepileptic, sedative-hypnotic and antiparkinsonism drugs involved benzodiazepines (46%, 3,725 hospitalisations, 14 hospitalisations per 100,000 people).

The rate of non-opioid analgesic-related hospitalisations fluctuated between 2003-03 and 2021-22, peaking at 39

11

Drug-Related Hospitalisations by Jurisdiction

The below sections describe trends in drug-related hospitalisations for each jurisdiction from 2002-03 to 2021-22. We

Figure 25. Age-standardised rate per 100,000 people of drug -related hospitalisations, by sex, Australian Capital Territory , 2002-03 to 2021-22.

Figure 26. Age92 re W* n BT /F4 9.96 Tf 1 0 0 1 84.384 496.75 Tm 0.38 0.122 0.451 rg 0.g T /F4 9.96 Tf 1 0 0 1 36 496.75 T

Figure 27. Age-standardised rate per 100,000 people of drug -related hospitalisations, by drug identified in the principal diagnosis, Australian Capital Territory , 2002-03 to 2021-22.

Note: Age-standardised rates were not calculated if the number of hospitalisations was less than or equal to 10 (please refer to our methods

Figure 32. Age-standardised rate per 100,000 people of drug-related hospitalisations, by sex, Northern Territory , 2002-03 to 2021-22.

Figure 33. Age-standardised rate per 100,000 people of drug-related hospitalisations, by remoteness,

Tre

Figure 36. Age-standardised rate per 100,000 people of drug -related hospitalisations, by sex, Queensland, 2002-03 to 2021-22.

Figure 37. Age-standardised rate per 100,000 people of drug -related hospitalisations, by remoteness, Queensland, 2018-19 to 2021-22.

Note: The size (area) of the bubble is proportional to the number of hospitalisations. In Queensland, data by remoteness area are only available from 2018-19.

Figure 40. Age-standardised rate per 100,000 people of drug -related hospitalisations, by sex, South Australia, 2002-03 to 2021-22.

Figure 41. Age-standardised rate per 100,000 people of drug -

Figure 44. Age-standardised rate per 100,000 people of drug -related hospitalisations, by sex, Tasmania, 2002-03 to 2021-22.

Provision of Tasmanian data between 200809 and 2015-16 was limited to drug related hospitalisations based on selected drug

Figure 48. Age-standardised rate per 100,000 people of drug -related hospitalisations, by sex, Victoria , 2002-03 to 2021-22.

Note: From 1st July 2011 to 30th June 2013 (i.e., between 2011-12 and 2012-13), there was a large decrease in public hospitalisations U H S R U W H G I R U W K H 9 L @ y.¿ ñ ’7Ec ñ p r r7`ä 0 o v Ef`ä qP0 R U Ef` R U•

Figure 52. Age-standardised rate per 100,000 people of drug-related hospitalisations, by sex, Western Australia, 2002-03 to 2021-22.

Figure 53. Age-standardised rate per 100,000 people of drug-related hospitalisations, by remoteness, Western Australia, 2012-13 to 2021-22.

Note: The size (area) of the bubble is proportional to the number of hospitalisations. Data on remoteness are only available from 2012-13.

Figure 54. Age-standardised rate per 100,000 people of drug-related hospitalisations, by principal diagnosis of mental and behavioural disorder due to substance use (A) and external cause of poisoning (B), Western Australia, 2002-03 to 2021-22.



Figure 55. Age-standardised rate per 100,000 people of drug-related hospitalisations, by drug identified in the principal diagnosis, Western Australia, 2002-03 to 2021-22.

Note: Age-standardised rates were not calculated if the number of hospitalisations was less than or equal to 10 (please refer to our methods document for details). Suppressed data are visible as gaps in the data series.