


Population ageing and apparent decline in initiation of injecting drug use in Australia

Olivia Price¹, Dan Lewer^{2,3}, Amy Peacock^{1,4}, Lisa Maher⁵, Farzaneh Zolalā⁶, Louisa Degenhardt¹, Sarah Larney^{6,7}, Seraina Agramunt⁸, Paul Dietze^{8,9}, Samantha Colledge-Frisby⁸, Nick Scott⁹, Joanna Wilson⁹, Rachel Sutherland




Objective

Among people who inject drugs in Australia, we examined temporal trends of **age**, **age at injecting drug use initiation**, **duration of injecting drug use**, and **year of injecting drug use initiation**. Based on Australian population size estimates of people who inject drugs, we **modelled the number of people injecting drugs for the first time** each year.


Data source

Twenty years (2000-2019) of repeat cross-sectional data from two national surveys with people who inject drugs in Australia - the **Illicit Drug Reporting System (IDRS)** and the **Australian National Syringe Program Survey (ANSPS)**.


- ‡ The **median age** of IDRS and ANSP participants has **increased** over the survey period (left)
- ‡ The **age at injecting drug use initiation** has **increased** with calendar time (middle)
- ‡ **Time since injecting drug use initiation** has **increased** over the survey period (right)



Modelled number of people injecting drugs for the first time in Australia



There was a **sharp decline** in the **number of people injecting drugs for the first time** in the **mid 1990s**.



The **peak in initiating injecting drug use** prior to this decline was **sensitive to model assumptions**, ranging from the **early 1980s to the early 1990s**.

Implications

1. People who inject drugs in Australia are an **ageing population**.
2. It appears this is driven by a **sharp decline in the number of injecting drugs for the first time in the mid 1990s**, and a **cohort of people who began injecting in the 1980s or 90s and continue to inject**.
3. Increasing age and injecting drug use duration have **implications for health service delivery**, as they may be accompanied by an **increase in chronic health conditions** and **injecting-related injuries and diseases**.

Author affiliations

¹National Drug and Alcohol Research Centre, UNSW Sydney, Sydney, Australia; ²Department of Epidemiology and Public Health, University College London, London, UK; ³Bradford Institute for Health Research, Bradford Teaching Hospitals NHS Foundation Trust, United Kingdom; ⁴School of Psychological Sciences, University of Tasmania, Hobart, Australia; ⁵Kirby Institute, UNSW Sydney, Sydney, Australia; ⁶Department of Family Medicine and Emergency Medicine, Université de Montréal, Montreal, Canada; ⁷Centre de Recherche du CHUM, Montreal, Canada; ⁸National Drug Research Institute, Perth, Australia; ⁹Disease Elimination, Burnet Institute, Melbourne, Australia



For further information, please contact Olivia Price o.price@unsw.edu.au