



MATHEMATICS ENRICHMENT CLUB.
Problem Sheet 18, September 24, 2018

1. A chess board is an 8×8 grid of squares coloured white or black so that no two adjacent squares are the same colour. Given tiles that are 2×1 grid squares, it is possible to cover the chessboard completely, and it takes precisely 32 tiles. Show that it is impossible to cover a chessboard with opposite corners removed.
2. Find all 3 digit numbers which are equal to the sum of the factorials of their digits.
3. In the diagram below, ABC is a circle of radius R with 3 tear-drop shapes inside. Each of the arcs $AC^{\circ}A^{\circ}$, $BA^{\circ}B^{\circ}$ and $CB^{\circ}C^{\circ}$ are of circles of the same radius, r . Find the area of each tear drop in terms of r .



5. The point M lies on the circumcircle of the equilateral triangle $\triangle ABC$, as shown in the diagram.



Prove that $AM = MB + MC$.

Senior Questions

1. Let f and g be real-valued, continuous functions defined on $-1 \leq x \leq 1$. We define the *inner product* of f and g , $\langle f; g \rangle$, as

$$\langle f; g \rangle = \int_{-1}^1 f(x)g(x) dx.$$

Consider the polynomials $p_0(x) = 1$ and $p_1(x) = x$.

- Two functions, f and g , are said to be *orthogonal* if $\langle f; g \rangle = 0$. Show that p_0 and p_1 are orthogonal.
- A function f is said to be *normalized* if $\langle f; f \rangle = 1$. Find factors α_0 and α_1 such that the polynomials $q_0 = \alpha_0 p_0$ and $q_1 = \alpha_1 p_1$ are not only orthogonal but also normalized.
- A set of polynomials that are all normalized and mutually orthogonal is called