

Solution Sheet 6, June 4, 2012

Answers

1. 6
2. only c and d are always true.
3. $x = 170; y = 13$ and $x = 170; y = 3$
4. $5\sqrt{2}$
5. Notice that using only the rules 1 and 2 ($(2x; y)$ and $(x; 2y)$ resp.) we can obtain all points of the form $(2^n; 2^m)$ and $\gcd(2^n; 2^m) = 2^{\min\{n, m\}}$: a power of 2. Furthermore, the operations $(x - y; y)$ and $(x; y - x)$ (as used in Euclid's algorithm), preserve the gcd. Hence points with a gcd that is not a power of 2 cannot be reached.
Conversely, these are the only points that can be reached. If $\gcd(a; b) = 2^m$, then $a = 2^m a'; b = 2^m b'$ with $\gcd(a'; b') = 1$. The point $(a; b)$ can be reached from $(a'; b')$ using rules 1 and 2 (apply each m and n times resp.).
Assume $a' < b'$. Since both $a'; b'$ are odd, $a' + b'$ is even, and can be reached from the point $(a'; \frac{a' + b'}{2})$. Notice that this point is closer to $(1; 1)$ than $(a'; b')$ was.
Continue this process until $a' = b'$, since $\gcd(a'; b') = 1$, this point is $(1; 1)$.