

MATHEMATICS ENRICHMENT CLUB.
Solution Sheet 1, April 30, 2016

1. Since any power of a number ending in the digit 6 is a number that ends with the digit 6, and $2^4 = 16$, the last digit of 2^{468} is 6.
2. There are 89 possible ways for Shaun to take the stairs. He can take different combinations of 1 and 2 steps, and each combination has a number of orderings as to when the 1 step or 2 steps are taken. For example, one combination is 6 1 step and 2 2 steps. The total number of ways to order the 1's and 2's is $8!$, but we don't care about the $6!$ ways to order within the 1's, and the $2!$ ways to order within the 2's. Hence this combination contributes

$$\frac{8!}{6! \cdot 2!};$$

ways for Shaun to take the stairs.

By completing the squares, one has $x^2 + 4x + 5 = (x + 2)^2 + 1^2$ and $x^2 + 2x + 5 = (x + 1)^2 + 2^2$. The former equation is the squared distance between the point $(x; 0)$ and $(-2; 1)$, while the latter is the squared distance between $(x; 0)$ and $(-1; 2)$.

Therefore, if we set let $P = (x; 0)$, $A = (-2; 1)$ and $B = (-1; 2)$, then we can think of S

Hence, $\sum_{i=1}^{100} n_i = 50$. Therefore,

$$\sum_{i=1}^{100} (n_i + 2)^2 = \sum_{i=1}^{100} n_i^2 + 4 \sum_{i=1}^{100} n_i + 400 = 200.$$

6. This is not possible.

Senior Questions

1. We introduce modular arithmetic for this solution; for example $p = r \pmod{5}$ means the remainder of p divided by 5 is r (also see https://en.wikipedia.org/wiki/Modular_arithmetic).

Trying a few prime numbers $p > 5$, one sees that $4^p + p^4$ is divisible by 5; that is $4^p + p^4 = 0 \pmod{5}$. We claim that this holds for every $p > 5$.

Note that $p = 1; 3; 7; 9 \pmod{10}$ covers all prime numbers. Moreover, if $p = 1; 9 \pmod{10}$, then $p^2 = 1 \pmod{10}$, which implies $p^4 = 1 \pmod{10}$. Similarly, if $p = 3; 7 \pmod{10}$, then $p^4 = 1 \pmod{10}$. In particular, for any prime number $p > 5$, one has $p^4 = 1 \pmod{5}$.

Also, it is easy to show by mathematical induction that $4^n = 4 \pmod{5}$ for all odd numbers n . Thus, $4^p = 4 \pmod{5}$ for all prime $p > 5$.

We conclude that $4^p + p^4 = 1 + 4 \pmod{5} = 0 \pmod{5}$, this proves our claim.

2. Let $v_A; v_B$ and v_C denote the velocity of Alex, Ben and Christ respectively. By the triangle's inequality, one has $AB + BC > AC$. Moreover, Alex and Ben both start at A and reach C at the same time. Hence $v_A > v_B$. Similarly, $v_A > v_C$.

Suppose Ben and Christ meets at the point O . Consider the interval of time in which Alex travels from point B to C : Since Ben meets Christ at O in this time interval, we can add another person Dean whom travels a distance of BO at constant velocity

and setting $y = x^3$ then using (3)

$$f(x^4) = xf(x) + x^3 f(x^3) = (x + x^5 + 2x^6)f(x): \quad (4)$$

Combining (2) and (4) yields

$$4x^3 f(x) = (x + x^4 + 2x^6)f(x):$$

Therefore, $f(x) = 0$ unless

$$4x^3 = x + x^4 + 2x^6: \quad (5)$$

But using (2)

$$f(x^5) = xf(x) + x^4 f(x^4) = (x + 4x^7)f(x);$$

and using (1) and (3)

$$f(x^5) = x^2 f(x^2) + x^3 f(x^3) = (2x^3 + x^4 + 2x^6)f(x):$$

Therefore, by similar arguments as before, $f(x) = 0$ unless

$$x + 4x^7 = 2x^3 + x^4 + 2x^6: \quad (6)$$

One can check that (5) and (6) can not occur simultaneously on the interval $(0;1)$. Thus $f(x) = 0$ for all $x \in (0;1)$.