SCHOOL OF BIOTECHNOLOGY AND BIOMOLECULAR SCIENCES

MICROBIOLOGY 1 (MICR2011)

SUBJECT GUIDE AND LABORATORY MANUAL

2022

Student I.D.

Course Information	
Course Code	MICR2011
Course Name	Microbiology 1
Academic Unit	School of Biotechnology and Biomolecular Sciences
Level of Course	

Graduate Attributes Developed in this Course		
Science Graduate Attributes	Degree of development *0 = no development* *1 = minimal* *2 = minor* *3 = major*	Activities / Assessment
Research, inquiry and analytical thinking abilities	3	Guided laboratory practicals, independent and collaborative lab research, assessment of open-ended investigations.
Creativity and motivation for intellectual development	3	Semester project and associat

Course Schedule					
Week	Date	Lecture		Practical	Tutorial
Week 1	Feb 14th	Introduction to MICR2011	JT	Aseptic technique	Sterilisation
	Feb 17th	Unicellular structure and function	JT		
	Feb 18th	Microbial phylogeny	JT		
Week 2	Feb 21st	Course Project	GL	Project DNA extraction	Phylogeny and classification
	Feb 24th	Microbial bioinformatics	MT		
	Feb 25th	Microbial metabolism 1	BB		
Week 3	Feb 28th	Microbial metabolism 2	BB	Biochemical techniques for characterisation	Biochemical techniques for characterisation
	Mar 3rd	Microbial evolution			

Assessment	
Assessment type	

Administration Matters
Expectations of Students

Additional support for students
x The Current Students Gateway: https://student.unsw.edu.au/ x Academic Skills and Support: https://student.unsw.edu.au/academic-skills x Student Wellbeing, Health and Safety: https://student.unsw.edu.au/wellbeing x Disability Support Services: https://student.unsw.edu.au/disability-services x UNSW IT Service Centre: https://www.it.unsw.edu.au/students/index.html



Contents:

WEEK 1 RISK ASSESSMENT 17
WEEK 1 LABORATORY CLASS 18
1. Welcome! 18
1.1 Allocation of places 18
1.2 Introductions 18
1.3 Laboratory safety 18
Microbiology Laboratory Safety Rules 19
Risk management 20
Risk Management Form 22
2. Media for the cultivation of microbes 26
2.1 Understanding media recipe composition 26
2.2 Aseptic technique 26
2.3 Making and sterilising liquid broth and agar plates 28
i) Luria broth 28
Materials 28
Procedure 28
ii) Pouring Luria agar plates 7T

WEEK 1 LABORATORY CLASS

1. Welcome!

1.1 Allocation of places

Students should proceed to the appropriate laboratory (Lab 11 or 12 in Biosciences South E26 as advised) where practical class groups will be organised.

1.2 Introductions

Students and tutors should get acquainted:

- Briefly introduce individual backgrounds.
- A brief discussion of course objectives and requirements.
- Explanation of assessable components of this course.
- Brief overview of laboratory and tutorial programs.

1.3 Laboratory safety

Note: In addition to reading and signing off on the material below all students will be required to complete the laboratory BABS Health and Safety Quiz in Moodle before next week's class (Week 2). You will not be able to proceed in further practical classes until you have completed this quiz.

In the laboratory, individuals are exposed to hazards not normally found in a regular classroom. It is essential that students follow all laboratory rules and regulations. Failure to follow established rules may result in dismissal of the individual from the class. Laboratories have certain standard safety equipment. These typically include:

- general-purpose fire extinguisher
- fire blanket
- eyewash
- safety shower
- cut-off switches for electrical and gas outlets.

It is the responsibility of the student to locate and know how to use the general safety equipment in the laboratory. Additionally, students should be aware of exits from the room in case of emergency, how to summon Campus Security (9385 6666), and how to obtain emergency medical assistance.

The microbiology lab has some additional safety considerations. Since individuals work with potentially pathogenic organisms care must be taken to prevent possible infection

I have read and understand the above rules and agree to follow them.

Signed_____Date_____

Name _____

Risk management:

For every task we perform in the laboratory that involves new equipment or substances that are potentially hazardous we are legally required to prepare a risk management plan (Workplace Health and Safety Act and Regulation 2011).

What is Risk Management?

It is about identifying hazards, assessing them and controlling the risk of the hazards. When identifying hazards consider where, how and who is carrying out work. The level of risk is assessed and used to prioritise implementation of risk control measures.

The risk control measures are:

- Eliminate the hazard(s)
- Substitute for something less hazardous
- Isolate the hazard from people
- Engineering controls e.g. plant guarding, work in fume cupboard or biosafety cabinet
- Administrative controls e.g. training, safe work procedures, signage
- Personal protective equipment (PPE) e.g. lab coat, safety glasses, gloves

To prepare a risk management plan/assessment you need to do the following:

- Identify Hazards: Source of potential harm to people, property or environment
- Assess Risks: Chance of something happening - consequence vs likelihood
- Eliminate or Control Risks

For the practical based studies you will undertaking in this course you will be required to read and sign the risk management plan associated with each individual laboratory class to provide evidence you have understood the associated risks and the control measures that have been put in place to minimise these risks. The risk management plans have been prepared by experienced staff however to help you understand this process we will ask you to identify the hazards and risks you may encounter in a microbiology lab and control measures we can use to minimise these risks and then discuss this with your fellow students and demonstrator.

An example of a risk management form for working with a microscope is shown on the next page. Note all risk assessments at UNSW are now submitted on-line using the

SafeSys database (http://safety.unsw.edu.au/node/93/).

Hazards and Risks	
Hazard Category	Ergonomic
Associated Harm	Back, neck or eye strain due to extended use or too high intensity of light source
Existing Controls	4 hour maximum booking allowed. Adjustable chairs provided. Training and risk assessment
Risk Consequence	Minor
Risk Likelihood	Unlikely
Risk Rating	Low
Additional Controls	-
Hazard Task	Exposure to biological materials
Hazard Category	Biologicals
Associated Harm	Infection
Existing Controls	PPE (lab

Describe hazards and risks associated with working with unidentified microorganisms.

Describe control measures that can be put in place to mitigate these risks.



maintain them free of external contamination is a basic microbiological technique. In this exercise you will have the opportunity of practicing sub-culturing and streak-plating techniques and of isolating organisms in pure culture. It all revolves around something we call aseptic technique. Accidental contamination may ruin your results completely.

Aseptic technique is largely a matter of common sense, but it is essential to realise that bacteria and fungi are present almost everywhere. Correct methods of handling cultures and apparatus must be perfected.

Consider carefully and remember the following points:

- Air contains many bacterial and fungal cells carried on dust particles or in water droplets.
- Any surface exposed to air quickly becomes contaminated, and if material is to be kept sterile exposure to air must be minimized.
- Instruments which can be sterilised by heating in a Bunsen flame (e.g. metal inoculating loops) can be left exposed, but they must be flamed thoroughly before use, and again before being replaced in the holder.
- Items of equipment that cannot be treated in this way are sterilised in wrappings or containers from which they must not be removed until actually needed. They must not be allowed to touch non-sterile surfaces during use. Plugs and caps of tubes and bottles must not be laid on the bench nor must sterile containers be left open to collect falling dust.
- Clothes, hair, skin and breath all carry a heavy microbial load and where strict asepsis (sterility) is essential (eg. surgery) sterilised gowns, caps, gloves and masks are worn. Even in normal microbiological work care must be taken to prevent contamination from the above-mentioned sources.
- A clean laboratory is the foundation of good aseptic technique.
- Microbial contamination in the lab is most often due to currents of non-sterile air. Ideally biosafety cabinets are used to minimize contamination. In the absence of a biosafety cabinet, a Bunsen burner can be used to create an updraft, thereby preventing contamination by settling dust particles. This protection can also be supplemented by keeping all windows and doors shut and by cutting down movement within the laboratory.
- Before commencing any operation, all necessary materials should be assembled on the bench for convenient access with provision for protecting sterile objects until needed.

At this stage of this practical class each tutor group will be divided into four mini-groups (with 4 students each). You will also use these mini-groups for the main course project.

Each group will initially be allocated to one of the following exercises:

1. Pouring agar plates,
2. Making Luria broths,
3. Or choosing your environment for the study of its indigenous microbes

Each group will then rotate through each exercise. It is anticipated that each exercise will take roughly half an hour.

2.3 Making and sterilising liquid broth and agar plates

Luria broth, originally know as Lysogeny broth, was devised by Giuseppe Bertani in 1952 for growing pathogenic *Shigella* lineages. It is now commonly used for the growth of enteric species such as the research workhorse *Escherichia coli* and in recombinant DNA work. It is a complex media that is easy to prepare and contains a wide range of nutrients with the following ingredients:

- Tryptone: peptides and peptones providing essential amino acids
- Yeast extract: vitamins, trace elements and certain organic compounds
- NaCl: sodium ions for transport and osmotic balance.

i) Luria broth

Materials:

- Tryptone
- Yeast extract
- Sodium chloride
- 50ml polyproylene centrifuge tube
- Spatula, weighing boat, balance
- Specimen container and pipettes.

Procedure:

1. Using a spatula, weighing boat and a balance, each student should weigh out the following ingredients into a conical (Erlenmeyer) flask. Rinse and dry the spatula between different chemicals to avoid mixing (no double dipping).
   - Tryptone: 0.2 g
   - Yeast extract: 0.15 g
   - NaCl: 0.2 g

2. Dissolve in 20 mL of distilled water.

3. Using a pipette aliquot 2 x 10 mL volumes into the 50 ml tube provided. Label the lid of your bottles with your name and date and place in the tray provided. The tubes will er1 (i)1 .9 (av)-(k)-1 (.)1TJ0 Tc 0 Tw 14.845 0 315 0

Is Luria broth defined or complex? Name one of the carbon and energy sources in the media.

For the three individual components state whether they are defined or undefined.

Tryptone:

Yeast extract:

NaCl:

ii) Pouring Luria agar plates

Materials:

- 40 mL Luria agar
- sterile petri dishes

Each student will be provided with 40 mL of molten Luria agar. This media has the same ingredients as the Luria broth above but 1.5% (weight/volume) agar has been added as a solidifying agent. Agar dissolves at 100ęC and so will dissolve during autoclaving (121ęC). The agar will not set until the temperature drops to below 40ęC. Agar can be kept in a 60ęC water bath to keep it molten before being poured into petri dishes.

Procedure:

1. Collect two petri dishes – take care when handling them as these dishes are sterile inside. Label the bottom of each dish with your name, tutor group and date. Place

What weight of agar (in grams) would be added to 40 mL Luria broth to give a 1.5% (w/v) concentration?

How do you think petri dishes are sterilised?

Why do we label the bottom of the petri dish?

3. MICR2011 course project

Whilst each weekly practical class involves independent laboratory activities, one exercise runs throughout the session and forms the basis of a total of 24% of the course assessment. In essence, the course project makes a comparison between traditional cultivation based microbiology and contemporary molecular microbiology. In your mini-group of four students will:

1) select an environment of interest
2) extract DNA from the environment of interest
3) identify microbes in that environment from DNA sequence data
4) isolate microbes from the environment of interest l2 -0 0 an env



WEEK 2
LABORATORY

OVERVIEW

This week we begin the course project. We will take two approaches to studying the microbial communities (microbiomes) of the samples you have chosen. The first is a contemporary molecular approach and the second is a traditional cultivation approach. The molecular approach involves extracting DNA directly from the sample and using DNA sequencing technology to determine what microorganisms are present. We will perform the DNA extraction today and analyse sequence data in subsequent weeks. The cultivation approach involves growing individual microbial lineages present in the sample and subsequently characterising them using biochemical tests, microscopy, and DNA sequencing. We will inoculate growth media today and do some cell staining and microscopy training exercises.

LEARNING OBJECTIVES

- Understand the difference between modern DNA sequencing-based techniques and traditional cultivation-based techniques for identification microorganisms.
- Understand the advantages and disadvantages of each approach.
- Develop skills for visualising microbial samples by light microscopy.
- Be able to Gram stain a bacterial sample and understand the basis of the stain.

ACTIVITIES

1. DNA extraction of project samples for sequencing.
2. Staining bacteria for observation by light microscopy. 2.-

Contents:

WEEK 2 RISK ASSESSMENT 35
WEEK 2 LABORATORY CLASS 36
1. DNA extraction of project samples for sequencing 36
Procedure 36
Materials 37
Step One - Sample collection and preparation 38
Solid samples (soil, food etc.) 38
Aqueous samples (seawater, pond water etc.) 38
Surface swab samples (skin, ear, benches etc.) 39
Step Two – Lysis 39
Step Three - Clean up (solid samples only) 39
Step Four - DNA binding and column washing 39
Step Five – Elution 40
Sequencing 40
2. Introduction to light microscopy 41
Parts of a light microscope 41
Setting up the light microscope 42
Resolution 44
Oil immersion 45
Microscopy trouble shooting 45
3. Staining bacteria for observation by light microscopy 46
Materials 47
Procedure 47
Smear preparation 47
Gram staining your heat fixed bacterial smear 49
Observations 49
4. Inoculating agar plates to obtain pure cultures and liquid broths as enrichment cultures 50
Materials 50
4.1 Procedure for inoculating samples to obtain single colonies (pure cultures) 50
4.2 Procedure for setting up enrichment cultures 50
Technique for subculturing agar plates for single colonies 51

WEEK 2 LABORATORY CLASS

1. DNA extraction of project samples for sequencing

DNA extraction underpins many modern techniques for the identification of microorganisms. Whether from a pure culture of a bacterial pathogen or a complex microbial consortium such as a stool sample, knowing what microbes you are dealing with starts with DNA extraction. DNA extraction from any given sample involves breaking open the cells, separa-3 (i)0.9-3 (ng w)- -(D)1 (her)-3.1 (f)1 (f)-
ttdebr (epar)-3 ar((epaf)1 (ef)-3 br (epaan)-1 (r)-and(ng opp-3.1 ((oder)er)-3 (p)- ()f)-3 (r0.9 nd om)2



Materials:

- PureLink Microbiome DNA purification kit
- Forceps, blades, swabs, saline solution, mortar and pestle
- Pipettes, tips, microfuge tubes, microcentrifuges and vortexes

Step One - Sample collection and preparation:

The method used to collect and prepare your sample will depend on the nature of the sample. Follow the appropriate guide for your sample type.

Solid samples (soil, food etc.)

i. Add 600 µL of Buffer S1 – lysis buffer to a bead tube (with red screw cap)
ii. Weigh out approximately 0.2 grams of your sample using the balance provided and add it to the bead tube. For food samples, it may be necessary to crush the sample using a mortar and pestle first.

Aqueous samples (seawater, pond water etc.)

i. Unscrew the blue nut on the top of the Vacuum Filtration Unit and remove the lid
ii. Remove a filter from its package using filter and place the filter in a container of deionized water to wet it completely.
iii. Remove the filter from the water using the forceps or soft-nosed tweezers and center it on a support screen. Make sure that the filter completely covers the screen.
iv. Using forceps, place a nitrocellulose filter disk onto the manifold (**A**) taking note of the position (1-12).
v. Replace the lid on the manifold and tighten the blue locking nut (**B**)
vi. Place stoppers in any positions around the manifold that do not contain filters. Add barrel extensions (max volume 50ml).
vii. Switch on the vacuum pump and filter the first 50ml of your sample, add more as the sample is filtered. Turn of the manifold and remove the lid (as in step i).
viii. Add 800 µL Buffer S1 – lysis buffer to a bead tube (with red screw cap) and add the filter membrane.



A: Setting the filter onto the manifold.
B: Filter loaded into the manifold.
C: Manifold with stoppers and barrel extensions connected to the vacuum pump.

Surface swab samples (skin, ear, benches etc.)

I. Add 600 µL of Buffer S1 – lysis buffer to a bead tube (with red screw cap)
II. Moisten the tip of a sterile cotton swab in sterile saline.
III. Swab the surface to be sampled thoroughly, making sure to use all sides of the swab.
IV. Place the tip of the swab into the bead tube and cut/break off the remainder. Ensure that any remaining handle does not prevent you from being able to cap the tube.

Step Two – Lysis

I. Label your bead tube with your group's identification on the side of the tube.
II. Add 100 µL of Buffer S2 – lysis enhancer to the bead tube. Cap tightly and mix tube contents briefly using the vortex mixer.
III. Incubate the tube in a heat block at 65°C for ten minutes.
IV. Homogenise the sample by shaking on a bead beater for 3 x 45 second cycles. *(Note: This step will be performed by the technical staff)*
V. Proceed to Step three (in the box below) if you are extracting from a solid sample. For all other samples proceed directly to Step Four.

Step Three - Clean up (solid samples only)

I. Centrifuge the tube containing your sample at 14,000 x *g* for 5 minutes.
II. Transfer up to 400 µL of supernatant to a clean microcentrifuge tube. Take care to avoid disturbing the pellet or transferring any debris.
III. Add 250 µL of Buffer S3 – Cleanup buffer to the supernatant and vortex immediately.
IV. Incubate on ice for ten minutes. Proceed to DNA binding (Step Four).

Step Four - DNA binding and column washing

I. Centrifuge the tube containing your sample at 14,000 x *g* for 2 minutes.
II. Transfer up to 500 µL of supernatant to a clean microcentrifuge tube, taking care to avoid disturbing the pellet or transferring any debris. Discard the tube containing the *pellet*.
III. Add 900 µL of Buffer S4 – B3 (ai)1 (eD0.003 Tw 0.28 0 Td[D)-3 (ube c)-1 (4di)1 (r)-3 (i)1 74 >>

Step Five – Elution

I. Transfer the column to a clean microcentrifuge tube labelled with your group identification.
II. Add 50 µL of Buffer S6 – Elution buffer to the column. Cap the column and allow it to sit at room temperature for one minute.
III. Centrifuge the column (in its microcentrifuge tube) at 14,000 x *g* for 1 minute. The DNA will now be eluted from the column into the tube, and is ready for quality analysis and sequencing. Hand your DNA extract to your demonstrator.

Sequencing:

The DNA extracted from your sample will be submitted to the Ramaciotti Centre for Genomics for further analysis. The quality of your DNA samples will be checked by gel electrophoresis and the quantity determined with a Nanodrop UV-

2. Introduction to light microscopy

The compound light microscope is an optical instrument designed for producing magnified images of objects using two or more glass lenses. The term light refers to the method by which light transmits the image to your eye. C*ompound* deals with the microscope having more than one lens. *Microscope* is the combination of two words; “micro” meaning small and “scope” meaning view.

The size of organisms and the internal structures to be observed determine the type of

Setting up the light microscope:



1. Switching on microscope

Turn the power switch on.

Rotate the voltage control clockwise to reduce light intensity or counter clockwise to increase intensity.



2. Specimen placement

Open the spring-loaded finger of the specimen holder and insert the slide.



3. Focus

Swing in the 10x objective .

Bring the specimen into focus using the course and fine adjustment knobs .



4. Interpupillary distance

Looking through the binocular tube, move the knurled dovetail slides until a suitable binocular vision is obtained.

5. Dioptre adjustment

Look at the image through the right eyepiece with your right eye, and focus on the specimen with the fine focus adjustment.

Looking at the image through the left eyepiece with your left eye, rotate the dioptre adjustment ring to focus on the specimen without using the focus adjustment knobs.

Your lens is dirty. Use lens paper, and only lens paper to carefully clean the objective and ocular lens. The ocular lens can be removed to clean the inside.

- Can't see anything under high power!
  Remember the steps, if you can't focus low power, you won't be able to focus anything under high power.

- Only half of the viewing field is lit, it looks like there's a half-moon in there!
  You probably don't have your objective fully clicked into place

Apparent fault	Possible cause	Correction
Field dark	Power (lamp) not on or turned down too low	Turn power on & check voltage
	Condenser diaphragm closed	Open diaphragm
	Lamp filament burnt out	Replace lamp
Colour of objects indistinct	Condenser diaphragm closed too far	Open diaphragm
Poor resolution	Condenser either too far open or too far closed	Adjust condenser diaphragm
Unable to focus on object	Cover-slip too thick	Replace
	Slide up-side down	Invert slide
	Focusing attempts too rapid	Use fine focus and adjust more slowly
	Objective has insufficient resolving power	Use higher power
	Objective covered with dried immersion oil from previous use	Clean with lens tissue and solvent

Materials:

- Three broth cultures, one pure culture of a Gram-positive organism, and one pure culture of a Gram-negative organism, and one unknown.
- Gram reagents.

Procedure:

The Gram stain is the most important differential stain in bacteriology. All bacteria can be classified as Gram positive or Gram negative, a characteristic that is produced by marked differences in cell wall structure and correlates with many other characteristics of the microorganism.

Smear preparation:

i. Obtain a clean glass microscope slide. Using an indelible marker such as a CHINAGRAPH £ pencil, draw a small circle on the slide where you intend to locate the smear.

ii. Sterilise loop by flaming in Bunsen burner as previously described (bacteriological loop method).

iii. Using aseptic technique, collect a portion of a single colony with the loop and mix with

Gram staining your heat fixed bacterial smear:

i. Flood air-dried, heat-fixed smear of cells for 1 minute with crystal violet staining reagent. Please note that the quality of the smear (too heavy or too light cell concentration) will affect the Gram Stain results.

ii. Wash slide in a gentle and indirect stream of water for 2 seconds.

iii. Flood slide with the mordant: Gram's iodine. Wait 1 minute.

iv. Wash slide in a gentle and indirect stream of water for 2 seconds.

v. Flood slide with decolorising agent (alcohol). Wait 15 seconds or add drop by drop to slide until decolorizing agent running from the slide runs clear.

fi (c)-1 (el)1 (-5 (w),-2 (as)-1 (ha1 (i)1 (2 (ai)0.9 (t)-30 Tc 0 Tw 1 0 v8)-3.1 (uns)-1 (cdTc 0 Tw
iii.o-I.

4. Inoculating agar plates to obtain pure cultures and liquid broths as enrichment cultures

Most techniques for the isolation of bacteria involve streaking an aqueous suspension of cells onto agar medium (to dilute the inoculum and obtain single colonies) and incubating for colony development. If a cell can grow on the media it will replicate until ultimately there is a colony of cells visible to the naked eye. The steps taken to obtain pure cultures are usually carried out a number of times in succession to safeguard against the transfer of contaminants. Sub-culturing is a useful microbiological technique and competence in it is desirable. Check your method with your demonstrator. If your plates are not adequate, repeat the initial isolation step.

In some instances, the organisms you are trying to detect may be present in your sample in very low numbers and may not grow successfully on plates directly from the original sample. In such instances the original sample can be incubated in a broth culture that will promote the growth of organisms – i.e. an enrichment culture. This is particularly relevant when trying to detect small numbers of possibly pathogenic organisms in human samples, especially blood cultures, or food samples or isolating microbes from environmental samples with specific abilities (e.g oil degrading bacteria from soil samples).

Materials:

- Samples of your choice for course project
- Luria broth - that you prepared in last week's class
- Agar plates:
  - Luria agar – plates you prepared in last week's class
  - LB20 – Luria agar with additional sodium chloride for marine samples
  - M9 – minimal salts agar with glucose as a carbon and energy source

4.1

Label the broths with your name, date and demonstrator's name and give them to your demonstrator. Suggest an incubation temperature

WEEK 3

LABORATORY

OVERVIEW

Isolation of microorganisms is required to unambiguously assign a property to the organism. Fulfilling Koch's postulates requires isolation. Phenotyping is a traditional means of identifying microbes in medical and food diagnostics. This week we will be checking out what grew on our plates from our samples, considering the purpose of enriched, selective and differential media, performing some phenotypic characterisation and learning how to count microorganisms.

S 0 Tw 10.0 P MCIDLBody13 BDC 0 gTT0 C2_0 0 12 164.7 65077 eq41 (eek)-1( )T2 56.7 611jEMC P 34.455ID

Contents:

WEEK 3 RISK ASSESSMENT.....55
WEEK 3 LABORATORY CLASS.....56
1. Examination of cultures obtained from project samples 56

WEEK 3 RISK ASSESSMENT

BABS Teaching Laboratory		MICR2011_RMF_Wk4
Student Risk Assessment		

Hazards	Risks	Controls
General	See below	**<u>At all times</u>** follow demonstrators' instructions when handling all biological substances, chemicals and equipment items. Notify demonstrator immediately of any spills or other incidents. Proper handwashing with antibacterial handwash before leaving the laboratory.
Biological agent (Microorganisms, Risk Group I and Risk Group II) Glass slides/coverslips Bunsen burner	infection cuts burns	PPE (lab coats, closed in shoes & gloves as required). Adhere to aseptic techniques. Work in PC2 laboratory or BSC if required. Dispose of any broken/used sharp/glass items immediately in Sharps containers provided. Never leave a lit Bunsen Burner unattended or reach over an exposed flame. Keep long hair tied back.
Chemical (hydrogen peroxidase & indole reagent, gram stain)	corrosive, irritant to eyes, respiratory system and skin	PPE (lab coats, closed in shoes, safety glasses & gloves. SDS.
Electrical Equipment (microscope, vortex)	Electric shock/ electrocution Exposure to biological agents (see above)	Avoid water/spillages when working with electrical

WEEK 3 LABORATORY CLASS

1. Examination of cultures obtained from project samples

Examine your plates from last week and select isolated colonies for further examination and subculture. Sub-culturing isolates will be a routine exercise until Week 8. If adequate biomass has formed on your plates or in your broth, Gram stain the colonies you have sub-cultured using the protocol described in last weeks' class.

Save and store photos of your plates. Record your Gram stain results here (you'll need these for your final report):

Suggestive learning activity:

Weekly group discussion for your course project - check Collaboration Checklist for suggestive topics

Don't forget to record group meeting minutes (meeting minute template) and develop weekly action plan (And stick to the plan!).

2. Selective and differential media

Most other organisms grow well but any haemolytic effect shown is irrelevant for differentiating between bacterial isolates, e.g. some strains of *Staphylococcus aureus* produce lysis while other strains show no lysis at all.

MacConkey agar (MAC)

Contains bile salts that inhibit many non -



What is the media component that makes Mannitol Salt Agar selective? Explain.

ÍCVcrÀÉU•d°Ì‡ÀÉU•d°Ì‡À”ÞÑÓ



Biochemical tests used for phenotypic characterisation

Staining bacteria and observation by light microscopy provides information about bacterial morphology, gram reaction, and potentially the presence of structures such as endospores. However, microscopic examination alone gives little information as to the genus and species of a particular bacterium.

To phenotype bacteria, some form biochemical analysis must be performed. The types of biochemical reactions each organism undergoes acts as a 'fingerprint' for its identification. Different species of bacteria have unique DNA and since DNA codes for protein synthesis, then different species of bacteria must, by way of their unique DNA, be able to synthesise different protein enzymes. Enzymes catalyse all the various chemical reactions of which the organism is capable. Therefore, different species of bacteria must carry out unique sets of biochemical reactions. By determining which enzymes are present in an unknown organism a description of that organism may be made and hence a step toward identification by phenotyping.

For example, all bacteria contain enzymes that can reduce oxygen to hydrogen peroxide and, in smaller amounts, to the more toxic free radical, superoxide. In most aerobes and facultative anaerobes, accumulation of these two toxic materials is prevented by the action of two enzymes, catalase and superoxide dismutase. Catalase is a hemoprotein enzyme of the oxidoreductase class that catalyses the conversion of hydrogen peroxide to water and oxygen. Superoxide dismutase is a copper-and zinc-containing enzyme present in all oxygen-using organisms that scavenges free radicals and converts them into H_2O_2 and O_2). Organisms that can tolerate exposure to oxygen always contain superoxide dismutase and may contain catalase. Lactic acid bacteria, being unable to synthesize heme molecules, do not contain catalase. Therefore, they are catalase negative. This group contains the enzyme peroxidase which catalyses the oxidation of organic compounds, such as $NADH_2$ by H_2O_2 which is thereby reduced to water. Lactic acid bacteria are therefore protected from H_2O_2 toxicity in the absence of catalase.
Enzymes are detected by the chemical reaction they carry out. However, it is necessary to have a means of observing and possibly measuring an enzyme's chemical reaction. One common measuring technique involves the reaction of a pH indicators that respond to the hydrogen ion, (H+), concentration by changing colour as a direct result of an enzyme's activity. The quantity of H+ in a solution is a measure of how acid or basic (alkaline) a solution is. Thus, if an enzymatic reaction produces acids, which increases the H+ concentration, pH-

Materials**:**

- Culture plate of *Staphylococcus*
- Culture plate of *Lactobacillus*
- Hydrogen peroxide

Procedure:

For each bacterial culture, place one drop of H_2O_2 (10%) on a glass slide.

Grow organisms for the test on any appropriate medium.

3.3 Indole

Amino acids may arise as a result of the hydrolysis of proteins, peptones and peptides. Tryptophan is an amino acid that can be used as an energy generating substrate by

3. Observe for the development of a blue colour within 3 minutes.

Results:

	Indole reaction
E. coli	
Proteus mirabilis	

An overnight broth culture of the yeast *Saccharomyces cerevisiae* (prepared by technical staff) will be counted using the spread plate method. Aliquots (100 µL) of appropriate dilutions will be spread evenly onto the surface of nutrient agar plates.

NOTE: Calculations have to be made to ensure that between 30-300 organisms are likely to be in one of the dilutions. You don't want more than 300 colonies on your agar plate because it gets too crowded to count. You don't want less because small numbers don't accurately reflect reality.

Materials**:**

- An overnight culture of *Saccharomyces cerevisiae* in Sabouraud broth (test sample)
- A series of suspensions of *S. cerevisiae* in Sabouraud broth (reference samples with known cell concentrations)
- 15 mL of sterile saline (per pair of students) will will

END OF WEEK 3



WEEK 4
LABORATORY

OVERVIEW

In the tutorial and lectures this week you studied microbially d (e)h

WEEK 4 RISK ASSESSMENT

BABS Teaching Laboratory		MICR2011_RMF_Wk7
Student Risk Assessment		

Hazards	

WEEK 4 LABORATORY CLASS

1. Winogradsky column

You have reviewed the Winogradsky column in this week's tutorial, but if you'd like some more information before answering the questions below, here's a little refresher:

The study of microbial ecology can be approached in 2 ways:

(i) By isolating, identifying, enumerating and studying the physiology of individual bacterial types of the ecosystem

(ii) By studying the activity (for example, chemical conversions) of the microbial population in the natural habitat.

One of the previous lab experiments, the exercise on isolation and identification of a given bacterial type, focuses on the former approach.

Bacteria and Archaea exhibit an astonishing metabolic diversity, which far exceeds that of the eukaryotes (animals, plants, fungi and other higher organisms). Bacteria and Archaea literally keep the biological world turning by recycling all the mineral elements necessary for life support. Two famous microbiologists pioneered the study of these processes, Sergius Winogradsky (1856-1953) and Martinus Willem Beijerinck (1851- - 0 . 0 0 1 e

not use water as the reductant; instead they use H_2S. The following simplified equations show the parallel.

$6\ CO_2 + 6\ H_20 \quad ÆC_6H_{12}O_6 + 6\ O_2$ (plant photosynthesis)

$6\ CO_2 + 6\ H_2S \quad ÆC_6H_{12}O_6 + 6\ S$ (bacterial anaerobic photosynthesis)

The purple sulphur bacteria typically have large cells and they deposit sulphur granules inside the cells. The green sulphur bacteria have smaller cells and typically deposit sulphur externally.

The sulphur (or sulphate formed from it) produced by the photosynthetic bacteria returns to the sediment where it can be recycled by *Desulfovibrio* - part of the sulphur cycle in natural waters.

Most of the water column above the photosynthetic bacteria is coloured bright red by a large population of purple non-sulphur bacteria. These include species of *Rhodopseudomonas*, *Rhodospirillum* and *Rhodomicrobium*.

These bacteria grow in anaerobic conditions, gaining their energy from light reactions but

1.1 Winogradsky column demonstration

At the front of the class there are two mature Winogradsky columns for you to observe. When indicated by your tutor, head to the front of the class and observe the different stratified communities present in the column. Unlike most occasions in microbiology, you can see the different microbial communities and differentiate some simply by their colouring.

In the space overleaf, you should sketch the column and indicate the different

Sketch the Winogradsky column below and indicate and microbial communities that you can identify with your group and tutor.

What is the purpose of the following ingredients that were added to the column?	
Mud	
Powdered cellulose	
Calcium sulphate	
Calcium carbonate	

Predict the location within the column and the chronological order of the

2. Soil based Microbial Fuel Cells

Harvesting energy from the environment is the primary activity of all living things. There are three principal ways to do this. Photosynthesis, fermentation and chemotrophic respiration. The latter is how we do it. As with humans, respiration in microbes involves the transfer of electrons to a suitable electron acceptor. For us this is oxygen, but for microbes there are many options. For example, bacteria can respire things like oxygen, nitrate, sulphate (seen in the Winogradsky column), iron, uranium, arsenic and even solid electrodes. Electrons can be transferred from bacterial cells to electrodes via soluble charge carrying molecules (electron shuttles), conductive appendages (nanowires or pili) or by direct cell contact.



The MudWatts provided (one per group) give us an opportunity not only to play with mud but also to bear witness to microbial respiration by linking the transfer of electrons from bacterial cells via an electrode to an LED. That's right, microbes can create enough potential in a bioelectrochemical system to make an LED blink. Not bad given the size of them. The following diagram illustrates the flow of electrons.





Materials:

- Carbon felt cathode and orange wire
- Carbon felt anode and green wire
- Vessel, hacker board, capacitor and LED
- Additional items including, multimeters, buzzers and clocks will be available in ensuing weeks to play around with.
- Mud – this is 'sampled' from the beautiful gardens around UNSW… discreetly. You wil be provided will either:
    1. Soil
    2. Soil supplemented with extra organic carbon (cellulose)
    3. Soil that has been autoclaved and is now sterile

Procedure:

Completely saturate your soil with water.

Bend both wires 90° where the plastic sheath ends. Straighten out the bare end of the wire. The green wire will be used to make the anode, and the orange wire will be used to make the cathode.

Insert the bare end of the anode (green) wire into the side of the thin felt disc. Try to keep the wire from exiting the felt. Repeat this step with the cathode (orange) wire and the thick felt disc.

Pack a layer of mud into the bottom of your container, at least 1 cm deep. Pat down the soil so that you have a smooth layer.

Place the anode (with the green wire) on top of the layer of mud.

Fill your container with more mud, at least 5 cm deep, pressing down firmly to squeeze out air bubbles. Let your mud rest for a few minutes and drain any excess liquid.

Place the cathode (with the orange wire) gently on top of the mud. Do not cover the cathode with mud.
Attach the hacker board into the indentation on the lid.
Pass the electrode wires through the lid.

Facing the semicircular indentation, the cathode (orange) should be on the left and the anode (green) on the right. Now press the lid down onto the jar to snap it into place.

Connect the cathode wire (orange) to the '+' and connect the anode wire (green) to the '-'.

Place the long end of the capacitor to pin 1 and the short end to pin 2. Connect the long end of the LED to pin 5 and the short end to pin 6.

It will take a few days for the microbes in the soil to colonise the anode at the bottom of the vessel and for the LED to start blinking. Over the next 2 weeks you will record the

group to take every 2-3 days. Record your results in the spreadsheet provided on the class OneNote Notebook (Content Library > Student Results > Soil microbial fuel cell results) and linked here.

3. Maintenance of project microbes

For those of you that haven't commenced or completed your phenotypic testing spend some time performing these tests as described previously. Don't forget to subculture your culture for future use. Photograph your culture plate and any of the tests you have performed. Keep your data and notes here.

Suggestive learning activity:

Weekly group discussion for your course project - check Collaboration Checklist for suggestive topics

Don't forget toTrBTer(c)-1 (er)-3 d( )5 ogroup(m)-3 (e(t)-3 (i-1 (gp(m)-3 (i-1 ((ut)-3.1 (es)]TJ0 Tc 0 Tw-

Number of colonies on lowest dilution (least dilute):

Number of colonies on intermediate dilution:

Number of colonies on highest dilution (most dilute):

Cell concentration in original solution (CFU/mL) =

Was the 10-fold dilution series apparent in your colony counts? If not, give reasons why:

END OF WEEK 4

WEEK 5 RISK ASSESSMENT

BABS Teaching Laboratory		MICR2011_RMF_COMP
Student Risk Assessment		

Workstation set-up	
Ergonomics	Musculoskeletal
Electrical	

WEEK 5 LABORATORY CLASS

1. Course project: sub-culturing, MALDI-TOF and storing project microbes

Examine your agar plates from last week. Take a photo and save to a location where you can access it for your final report (create a MICR2011_Report folder in your student drive). Based on colony morphology do you have a pure culture or mixed lineage growth on your plate? Pick a colony and subculture onto a new plate. Check next week that your microorganism is still alive. To ensure the culture is not lost during the following weeks stocks of the culture can be frozen and revived at a later time point if necessary as follows.

Procedure:

- Only continue today with the following if you are sure you have a pure culture. Otherwise, this step can be performed next week.
- Obtain a culture plate showing good growth of a pure culture of your isolate.
- Aliquot 0.5 mL of Luria broth (containing 15% glycerol) into a cryotube.
- Using a sterile loop, pick two or three colonies and transfer and suspend in the cryotube from above.
- Labeled with your name and date. Give this to your demonstrator who will store it at -80ęC until required.

Identify your microorganism

Once you are satisfied that you have a pure culture, you may begin your identification. You will use two methods: a more classical biochemical characterisation, and phylogenetic analysis of the 16S rRNA gene from your microorganism.

Flow charts of the tests required to identify your microorganism are provided in Appendix 1 and are adapted from Bergey's Manual of Determinative Bacteriology. The stains and tests listed (and instructions for performing the tests) are available in the lab and you should discuss with your tutor the best way to identify your microorganism. You will need to start with a Gram stain to determine the Gram reaction and cell morphology. This will also help you determine if you have a pure culture.

In the next practical, you will sequence part of the 16S rRNA gene from your microorganism and, using the skills you will learn today, identify your microorganism using phylogenetics.

MALDI-TOF mass spectrometry analysis of project isolates

Mass spectrometry (MS) is a relatively recent addition to the microbiologist's toolkit. MS can be used to create a "fingerprint" of the molecular weight of the most abundant proteins in the cell. These proteins are predominantly ribosomal proteins and there is sufficient variation in the molecular weight between genus, species, and sometimes strains to determine the microbe's identity.

The major advantage of using MS to identify microorganisms is the speed of identification. From a pure culture, you can identify the microorganism in less than an hour. Almost all MS instruments for bacterial identification use matrix-assisted laser desorption/ionization (MALDI) to vaporise and ionize your sample microorganism for MS. Your sample is mixed with a “matrix” solution, in this case we will use Ŭ-Cyano-4-hydroxycinnamic acid (HCCA) that will absorb energy from the UV laser of the MALDI to generate free radicals and heat that will ionize and vaporise the sample.

The ionised sample vapour containing proteins from your bacterial sample is then fed into a Time-Of-Flight (TOF) mass spectrometer, hence the name MALDI-TOF. The TOF spectrometer will measure the mass/charge ratio (m/z) of the proteins in the sample creating a fingerprint of the molecular weights of the most abundant proteins in the cell. To identify our microorganism we compare the spectra from the 2-20 kDa range of proteins against a database of spectra from previously characterised bacteria. In this way we can match the MALDI-



(indicated right) (Carbonelle *et al* Clinical Bioc emistry 2011)

Workflow used to analys
with the Bruker MALDI TOF Biotyper.

2. Introduction to sequence analysis

Bioinformatics now plays a central role in medical and environmental microbiology. From tracking the evolution of pathogenic viruses, to identifying a fungal isolate, or identifying bacteria in a sputum sample, making sense of DNA sequence data is crucial.

Over the past decade there has been a technological revolution in sequencing, with the costs dropping dramatically and the amount of sequence data increasing. For example in 2001 the cost of the Human Genome project was $100 million. Today genomes can be sequenced for around $1000.

This revolution has meant that sequence data is much more readily available to a wide range of scientists than was previously the case. In microbiology, this has resulted in the capacity to deeply sequence the DNA of microbes from any sample and readily identify community composition and function. As a consequence, much of the skill involved in medical and biotechnological research has moved from the “wet lab” to the computer lab, and you will need to become comfortable using (or even developing) many types of bioinformatic software during your career.

One good thing about playing around with bioinformatics is that you can’t really do any harm, spread a dangerous virus, burn down the lab, etc. You can probably take down a server with some bad code but this can generally be fixed pretty easily. Feel free to explore the software we will use (and any other similar packages you can find). Much newly developed software is open source, so if you are computationally inclined downloading, installing and trying out new methods is a great way of advancing your

knowledge. If you are less into computers then there are still many ways to get the information you need out of sequence data and move on.

There are many different ways to do most analyses, and as long as you develop the conceptual knowledge of what you do at each step of the process you have the flexibility to move between different tools as you see fit. Understanding and flexibility ensure you'll be able to deal with things when they don't go as expected and build checks into your workflows to ensure things stay on track.

We will use both old and new sequencing methods in this course. The newer methods are currently known as “next


V2
V3
V4
V5
V7
V9
R6: 1,301–1,542
R5: 1,0
R4: 751–1,050
R3: 501–750
R2: 251–500
R1: 1–250

The secondary structure of the SSU rRNA molecule is useful to understand the methods we use to explore diversity using the gene. When folded in two dimensions RNA forms a series of stems and loops. In general, as the stems require base-pair bonding, it is important that these regions of the gene are conserved. Nucleotide mutations in stem regions can stop secondary structure formation and render this essential molecule useless and the organism will die. That's a strong selection pressure for conservation. Loop sequences on the other hand don't base pair so they mutate more and are less conserved.

When you learn and read about using Illumina sequencing to examine microbial assemblages from different environments you will hear about variable regions V2-V3, or V4, or V6. These designations refer to specific sub-parts of the SSU gene that contain variability between species so are useful for telling one species from another.

Exercise 1. Accessing software for phylogenetic analysis of SSU rRNA sequences

Today you will be working in pairs and using a software package called MEGA for analysis of your mock SSU rRNA sequences. You will use MEGA to analyse your 16S rDNA Sanger sequencing results of your pet microbe for your final report.

The MEGA software package is freeware that allows us to visualise and manipulate sequence data, create multiple alignments, estimate evolutionary distance, and build phylogenetic trees, amongst many other uses. You can access MEGA using your student zID through myAccess:

https://www.myaccess.unsw.edu.au

Click on Access my apps and select MEGA

Opening an Alignment

The *Alignment Explorer* is the tool for building and editing multiple sequence alignments in *MEGA*.

1. Launch the *Alignment Explorer* by selecting the Align > Edit/Build Alignment on the launch bar of the main *MEGA* window.
2. Select Create New Alignment and click OK A dialog box will appear asking, “Are you building a DNA or Protein sequence alignment?” Click the button labeled “DNA”

3. From the *Alignment Explorer* main menu, select Data > Open > Retrieve sequences from File. Select the ‘ Micro2011_16S_rRNA_sequences.fasta’ file from the location that it was saved. You should see something like this:

Notice the different coloured blocks for each nucleotide (A= green, T= red, C = blue, G = purple)

Use the scroll bar at the bottom to see how long each of the sequences are.

You can manipulate the order of the sequences by dragging and dropping. Try moving the sequences up or down in order by dragging the names.

For example, move Candida_glabrata down so it is above the two Saccharomyces sequences at the bottom. Then move all the eukaryotes to the bottom (Saccharomyces, Candida and Penicillium)
Double click in the “Group Name” box next to one of the sequences and type in something to represent the Genus name. e.g. ‘Pro’ for *Prochlorococcus*. Do this for the rest of the groups (e.g. ‘Bord’ for the *Bordetella* sequences). Group *Candida* with the *Saccharomyces*. .

Aligning Sequences Using Muscle

Before you can do ANYTHING with your sequences they must be aligned (matched up so that you compare the right base pairs to each other).

1. On the *Alignment Explorer*

at the start and at the finish. This is a problem because you can't accurately compare sequences of different length. To resolve this we need to do one of two things:

- Delete the base pairs of all the other sequences from the start and end so they match up with NATL1. This will allow us to keep NATL1 in our tree but remove a lot of data we use to build the tree, or

- Remove NATL1 from the alignment before we make the tree, and just trim the ends to make the other sequences the same length.

Your decision will depend on whether you are interested in the position of NATL1 and the intergroup similarities in *Prochlorococcus*, or more so in the rest of the sequences?

If you want to remove NATL1 click on the name and then Edit > Delete to remove it.

OR

If you want to keep all your species for now, delete all the columns in the alignment from the start of the alignment to the start of the NATL1 sequence, and from the end of the alignment to the end of the NATL1 sequence, by highlighting the appropriate grey boxes at the top of the alignment and then Edit > Delete.



3. Select all five records again and display the full GenBank record (including sequences) by selecting Summary > GenBank (full) to load the data in the correct format. Once that has loaded, you can import these closely related sequences into MEGA using the Add to Alignment button at the top of the screen.



4. The related sequences will now be added to your Alignment Explorer window and you can repeat your alignment and phylogenetic tree construction (Exercises 2, 3, and 4) using your additional related SSU sequences. This will be particularly important for Week 11 when you have your unknown project samples and will need to identif-3 (E) b8-3 (12.9r)-TJ0 -1.15 TD[3,EEK 5)1gCer4 (at)-32 (pl)1 ps2ye yourt21 (er)-3 (i)1



WEEK 7
LABORATORY

OVERVIEW

Contents:

WEEK 7 RISK ASSESSMENT 108
WEEK 7 LABORATORY CLASS 109
1. DNA extraction from sub-cultured project isolates 109
Materials 109
Step One - Sample collection and preparation 109
Step Two - Lysis 109
Step Three - DNA binding and column washing 109
Step Four - Elution 110
2. Assessment of DNA extract quantity and quality 110
Materials 111
Procedure 111
Results 111
3. PCR amplification and sequencing of SSU rRNA genes 112
Materials 113
Procedure 113
4. Record the results of your Microbial Fuel Cell 103

WEEK 7 RISK ASSESSMENT

BABS Teaching Laboratory		MICR2011_RMF_Wk9
Student Risk Assessment		

Hazards	Risks	Controls
General	See below	**At all times** follow demonstrators instructions when

WEEK 7 LABORATORY CLASS

1. DNA extraction, biochemical characterisation, and antibiotic resistance analysis from sub-cultured project isolates

Working in your mini-group follow the protocol below to extract DNA from the microbes isolated by your group. We're using the same DNA extraction kit as we did previously so refer back to Week 2 to get an overview of the protocol. Detailed instructions are given below.

Materials:

- PureTc 0 Tw 3.961 0 Td(:)1ow( )Tj0 0.12-2.3 (i(:)1o)-TfE.9 (l)1 (o84)1oooK robes

13. After the second spin, transfer the column (with DNA now bound to the filter inside) to the clean collection tube provided.
14. Add 500 µL of Buffer S5 – Wash buffer to the column, cap it and centrifuge for 1 minute as above.
15. Discard the flow through, replace the column in the tube and centrifuge for an additional 1 minute (this step removes residual wash buffer from the column).

Step Four - Elution

16. Transfer the column to a clean microcentrifuge tube, labelled with your group identification and the date.
17. Add 50 µL of Buffer S6 – Elution buffer to the column. Cap the column and allow it to sit at room temperature for one minute.
18. Centrifuge the column (in its microcentrifuge tube) at 14,000 x *g* for 1 minute. The DNA will now be eluted from the column into the tube and is ready for quality analysis PCR and sequencing.

Continue - Biochemical characterisation of project isolate

You will continue with the biochemical characterisation of your project isolate using the guide provided in the Appendix.

methods for checking DNA quantity and purity that we won't pursue today include agarose gel electrophoresis combined with fluorescence imaging or quantifying directly with DNA binding fluorescent dyes using Qubit technology.

Materials:

- DNA sample from step 1 above
- Pipette and tips
- Nanodrop and Kimwipes™

Procedure:

1. To determine the absorbance of various wavelengths of electromagnetic radiation by DNA in elution buffer, first you need to see what the elution buffer looks like itself. Aliquot 2 µL of Buffer S6 – Elution buffer onto the Nanodrop stage, lower the arm and instruct the software to take a blank reading. This tells the Nanodrop that this is what the background of the sample looks like. Wipe the droplet off the stage with a Kimwipe™/tissue.

2. Now aliquot 2 εL of your DNA sample onto the Nanodrop stage and instruct the software to measure the DNA concentration. This will give you a spectrum and a concentration for your DNA.
3. Your spectrum should look something like this. The absorbance maxima (peak) of the spectrum should be at 260 nm. It is this reading that is used to calculate the DNA concentration. A 50 ng/µL solution of DNA will give you an absorbance at 260 nm of 1 so you can easily figure out DNA concentration from absorbance. The software also gives you absorbance readings at 230 nm and 280 nm. The ratio between the 260 nm and 230 nm readings (260/230 ratio) and the ratio between the 260 nm and the 280 nm readings (260/280 ratio) give you information on purity. The 260/280 ratio should be close to 1.8 for pure DNA. The 260/230 ratio should be above 1 if the sample is free from carbohydrates and phenolics. If your 260/230 and 260/280 ratios are not as expected you should not trust the concentration as measured by the Nanodrop. Note that the Nanodrop tends to slightly overestimate the DNA concentration.



Results:

Absorbance (230 nm):

Absorbance (260 nm):

Absorbance (280 nm):

260/230 ratio:

260:280 ratio:

DNA concentration (ng/µL):

Draw the absorbance spectra of your DNA sample below:



3. PCR amplification and sequencing of SSU rRNA genes

The DNA sequence that encodes the SSU rRNA sequence is used for taxonomical classification of microbes as there are conserved regions in the sequence (due to the

conserved function of SSU rRNA) and hypervariable regions that enable us to distinguish between close relatives. The gene encoding SSU rRNA is approximately 1500 base pairs long and can be amplified in a single PCR to obtain the sequence for the whole gene. Discuss with your demonstrator what primers should be used for sequencing.

END OF WEEK 7



12 5624 795E245.92Y10 12 5R24 20070770508 4085( )Tj0 0.1/ArtifPCID 1 >>B6C 0 0.47142.5 /

Contents:

WEEK 8 RISK ASSESSMENT 117
WEEK 8 LABORATORY CLASS 118
Materials 118
Procedure 118
Results 118
Phenotyping verses genotyping 120

WEEK 8 LABORATORY CLASS

Materials:

- x 4 plates nutrient agar (per bench group)
- x 4 plates mannitol salt agar (per bench group)
- x 4 plates Sabouraud agar (per bench group)
- x 4 plates MacConkey agar (per bench group)
- x 2 sterile "Stomacher" bags (per bench group)
- x 4 x 50 ml volumes physiological diluent (saline) (per group)
- x 30 g of an unprocessed food in a sterile petri dish (per group)
- x 30 g of a processed food in a sterile petri dish (per group)

Macroscopically examine the plates set up last week. Do you have any evidence of different population sizes on the processed and unprocessed foods?

Does comparative growth on different selective media and the colony appearance of the populations suggest any qualitative differences in the populations of the two foods?

Comment on the predominant colony types on each food	

Phenotyping verses genotyping:
DNA was isolated from three bacteria that grew from the food sources. You are provided with three 16S sequences from these bacteria. The sequences have been uploaded in fasta format to the MICR2011:Moodle page under Week 7.



WEEK 9
LABORATORY

OVERVIEW

Bacteriophages are viruses that infect and multiply within bacteria. Bacteriophages play a major role in regulating the abundance of bacteria in the environment, including the human body. Bacteriophages are used to prevent the souring of petroleum resources caused by sulphate reducing bacteria and to prevent, and even treat, bacterial infection. This week you will learn to quantify bacteriophage and examine bacteriophage mediated transfer of genes between bacteria in a process called transduction.

LEARNING OBJECTIVES

- Bacteriophages are obligate intracellular parasites and do not form independent colonies. This poses challenges for standard culture-based enumeration. Student's should understand how to cultivate and enumerate bacteriophages using plaque assays.
- Horizontal gene transfer allows bacteria to exchange DNA between species and accelerates bacterial evolution. Bacteriophages facilitate horizontal gene transfer through transduction and students should understand how this occurs and how we

Contents:

WEEK 9 RISK ASSESSMENT 112
WEEK 9 LABORATORY CLASS 113
1. Bacteriophage propagation and enumeration 113
Materials 114
Procedure 114
Dilution of the phage stock 114
Phage absorption and overlaying agar 114
2. Gene transfer by transduction 115
Materials 116
Procedure 116

WEEK 9 LABORATORY CLASS

1. Bacteriophage propagation and enumeration

Bacteriophage infection consists of a number of sequential events (refer to lecture by Prof. Peter White). The phage is first absorbed by the host cell, then replicates and assembles into mature

Materials:

Per mini group of students:

- 1 x 0.5 mL PRD1 bacteriophage stock

- 1 mL 24 h Z broth culture of *Salmonella typhimurium LT2*

- 10 mL Z broth
- 4 x Z agar plates (1 for control, 1 for each phage dilution to be plated) TjEMC /P 4MCID 34 >>E

Panel A shows a schematic of the structure of the bacteriophage P1, with head and tail

END OF WEEK 9

WEEK 10
LABORATORY

OVERVIEW

Today you will be looking at how we can track the spread of viruses (or many infectious agents) during an outbreak. Specifically, you'll be looking at a case study of Ebola infection and using ELISA analysis of blood serum samples to determine the source of the infection.

LEARNING OBJECTIVES

- Understand how viral pathogens can be detected and monitored in infected individuals using an ELISA.
- Understand how the combination of molecular detection techniques and patient meta-data is used determine the epidemiology of an infectious agent and trace the source of infection in an outbreak.

Contents:

WEEK 10 RISK ASSSSMENT 120
WEEK 10 LABORATORY CLASS 121
1. Background 121
Ebola virus 122
History 122
Transmission 123
Signs and Symptoms 124
Control 124
2. Epidemiological investigation 125
3. ELISA analysis of mock serum samples 127
Materials 128
Procedure 128
4. Tasks for the investigation team 129
Tasks .....

WEEK 10 LABORATORY CLASS

1. Background



Figure 1. Reported active human transmission in 2014-2016 (CDC website)
Background by Professor Vincent Racaniello, author of “This week in Virology”
Reference: http://www.microbe.tv/twiv/twiv-341/
To see more about Vincent: http://www.virology.ws/about/

Ebola virus:

The widespread outbreaks of Ebola virus in the West African nations, together with the fatality rate of up to 90% have propelled this virus into the limelight in recent years. What is this virus and where did it come from?

History:

Ebola virus was first identified in 1976 and is named after a river in the Democratic Republic of the Congo (formerly Zaire). Outbreaks have been sporadic since its discovery, and several strains have been identified in outbreaks across the African continent. Between 2014 and 2016 there was the largest Ebola outbreak in history, resulting in the most human fatalities since the discovery of the virus. Following the first confirmed case in

Outside of Africa there have been several cases of Ebola virus reported in countries

Group 3 remains in Freetown where they work with staff and officials at Abanita Hospital. The group includes:

- Prof Peter White
- Prof Zuri Babatunde
- Prof Leighton Scott
- Dr Jai Tree
- Mr Brandon Sy
- Dr Tendai Idowu

September 24th 2018

After collecting samples from the Abanita Hosptial in Freetown, the epidemiological research team travel to the MRC Unit in The Gambia to deliver samples of Rift Valley Virus and Ebola for sequencing. The experiments will take 3 weeks, so Brandon Sy and Dr Tendai Idowu leave Freetown early for the MRC Unit and Dr Jai Tree stays behind in Freetown to attend the Grand Opening of the WABIRT laboratories at the end of the week.

September 28th 2018

After observing and supporting the implementation of the new testing kits in Kenema and Kambia, the AusAID teams return to Freetown for the Grand Opening of the new Australian Foreign Aid funded WABIRT laboratories. Over 120 invited guests attend including those from WABIRT, University of Makeni, Njala University, the Sierra Leone Ministry of Health, the Australian embassy, the Sierra Leone government, a number of local hospitals and many other sources.

Both Dr Rowena Benton and Dr Jenny Morgan have had headaches and sore muscles for the previous 2 days. They do not attend the Opening and Dr Jenny Morgan is admitted to Abanita Hospital in the early afternoon with some blood in her vomit.

On examination she was unwell, with a high fever of 39.9°C and signs of bruising and blood from her eyes and nose. Antigen testing (using the new Australian antigen assay) of a blood sample reveals that Jenny had been infected with Ebola virus. Over the subsequent 12 hours Jenny's haemorrhagic symptoms worsens accompanied by seizures and loss of consciousness. From there, her condition deteriorates and she died at 11:45 pm the following day.

October 6th 2018

Seven people who attended The Grand Opening became ill with Ebola-like symptoms and are admitted to Abanita Hosptial. These include:

Prof Peter White
Dr Afia Ayodele
Prof Zuri Babatunde
Prof Subira Arendse
Dr Masamba Ejiro
Mr Amadi Afolabi
Dr Tinashe Kayode

Due to the death of Drzur anf Zm Td[P)1 (r)0 -1.15 ..9 (A)1 (f)-3 (ol)1 (abi)]T/emBDC /TT21 (i)14.45 -1

obtaining samples under your direction. The official may also help you to trace the whereabouts of specific individuals prior to the Opening. To aid your investigation you have been given a state-of-the-art Enzyme Linked ImmunoSorbent Assay or ELISA (developed at UNSW and commercialised through UNSW Innovations, the commercial arm of the University), which can detect with high sensitivity Ebola antigens (the Ebola virion) in blood samples.

Aim: To trace outbreak of Ebola virus, and identify the source of the outbreak

- First of all find how many of the seven attendees of the Grand Opening who became ill have the Ebola virus. You will also test sera from Drs Benton and Morgan. This will help you become familiar with the ELISA. Include a positive and negative control.

Suspected Cases	Ebola Test Result

Materials:

- Pipettes and tips
- Plates
- A positive test sample and a negative test sample
- Paper towels
- Antibody solution
- Washing solution
- Colour reagent solution (TMB)

On the 6th October blood samples are taken from those people in Freetown that came into contact with Drs Morgan and Benton, and attendees from The Opening. You have blodd samples from:

- Dr Jenny Morgan and Dr Rowena Benton
- 6 samples from the Kenema Team
- 10 samples from the Kambia Team
- 15 samples from the Abanita Hospital
- 7 people displaying symptoms after the Grand Opening
- 15 samples from passengers on flight KQ504 that were still in Freetown

- 120 samples from The Opening are not available for testing.
- 125 samples from flight KQ504 are also not available.

Procedure:

1. Label the tubes appropriately using the marker provided.
2. Your team will run an ELISA assay on your samples using the microtiter plates provided.

   Initially each team will do one positive and one negative control. Add 100 µl of a positive test solution into two wells and 100 µl of a negative test solution into a different set of two wells. Be sure to write down which wells are positive and which are negative controls. 100µl should fill one third of the well.
3. Add 100 µl of your nine suspected Ebola virus samples into each of two wells. Record which wells contain your samples.
4. After adding the samples leave the plate on the lab table undisturbed for 5 minutes.
5. Shake off the fluid into a nearby sink or designated container, making sure that the fluid has emptied from each well. Tap the plate upside down onto the paper towel to remove any excess liquid or bubbles.
6. Wash the plate using 150µl of washing solution in each well. Repeat a total of three times.
7. Remove excess solution and bubbles by again tapping the plate face down onto a paper towel.
8. Add 100 µl of the enzyme conjugated antibody solution to each well.
9. Allow for 5 minutes incubation time on the lab table and then shake off the fluid.
10. Repeat steps 6 and 7.
11. Add 100 µl of the colourising (TMB peroxidase substrate) solution to each well.

Gram Positive Cocci ID Flowchart






hart
Lactose Negative Flow C
Edwardsiella tarda
otuberculosis
Yersinia pseud
Edwardsiella tarda
Morganella morganii
+
Providencia stuartii
Indole Test